

DRUG DETERMINATION POLICY

Title: DDP-19 Benlysta

Effective Date: 09/21/2020



Physicians Health Plan
PHP Insurance Company
PHP Service Company

Important Information - Please Read Before Using This Policy

The following policy applies to health benefit plans administered by PHP and may not be covered by all PHP plans. Please refer to the member's benefit document for specific coverage information. If there is a difference between this general information and the member's benefit document, the member's benefit document will be used to determine coverage. For example, a member's benefit document may contain a specific exclusion related to a topic addressed in a coverage policy.

Benefit determinations for individual requests require consideration of:

1. The terms of the applicable benefit document in effect on the date of service.
2. Any applicable laws and regulations.
3. Any relevant collateral source materials including coverage policies.
4. The specific facts of the particular situation.

Contact PHP Customer Service to discuss plan benefits more specifically.

1.0 Policy:

This policy describes the determination process for coverage of specific drugs.

This policy does not guarantee or approve benefits. Coverage depends on the specific benefit plan. Drug Determination Policies are not recommendations for treatment and should not be used as treatment guidelines.

2.0 Background or Purpose:

Benlysta is a specialty drug indicated for Systemic Lupus Erythematosus (SLE) and is useful for specific organ system symptoms. These criteria were developed and implemented to ensure appropriate use for the specific symptoms detailed below.

3.0 Clinical Determination Guidelines:

Document the following with chart notes

A. Systemic Lupus Erythematosus (SLE).

1. Age: at least 18 years.
2. Diagnosis and severity.
 - a. Active moderate to severe systemic lupus erythematosus refractory or intolerant to other immunosuppressive drugs.
 - b. Autoantibody positive: ANA at or above 1:80 and/or anti-dsDNA at or above 30 Units/mL.
3. Chronic other therapies: contraindication, inadequate response or significant adverse effects to triple therapy listed under moderate or severe disease for four months (see Appendix I).
 - a. Moderate disease: prednisolone (≤ 7.5 mg per day); hydroxychloroquine; and azathioprine, methotrexate, mycophenolate mofetil OR cyclosporine.

- b. Severe disease: prednisolone (≥ 7.5 mg per day);hydroxychloroquine; and mycophenolate OR cyclosporine.
4. Dosage regimen:
- a. Adults: Benlysta intravenous (belimumab IV): 10mg per Kg every two weeks times three, then every four weeks or Benlysta SC (belimumab): 200mg every week.
 - b. Pediatrics: Benlysta intravenous 10mg per Kg every two weeks, then 10mg per kg every four weeks.
5. Approval.
- a. Initial: six months.
 - b. Re-approval: one year (decrease signs and symptoms of disease).
 - c. Administration: medication falls under site of care policy (see DDP-08).
6. Exclusions:
- a. Concurrent Disease: severe active lupus nephritis or central nervous system (CNS) lupus.
 - b. Concurrent Medications: other biologics or intravenous cyclophosphamide.

4.0 Coding:

AFFECTED CODES				
HCPSC Code	Brand Name	Generic Name	Billing Units (1U)	Prior Approval
J0490	Benlysta	belimumab	10mg	Y

5.0 References, Citations & Resources:

1. Lexicomp Online®, Lexi-Drugs®, Hudson, Ohio: Lexi-Comp, Inc.; Benlysta, accessed July 2020.
2. The British Society for Rheumatology guideline for the management of SLE in adults: Executive Summary. Rheumatology 2018;57.
3. DDP-08 Site of Care for Administration of Parenteral Specialty Medications.

6.0 Appendices:

See page 3.

7.0 Revision History:

Original Effective Date: 12/1/2011

Next Review Date: 07/22/2021

Revision Date	Reason for Revision
7/19	Annual review; replaced abbreviations
6/20	Annual review; removed mild disease other therapies replaced abbreviations, approved by P&T Committee 8/26/20.

Appendix I: SLE Treatment Strategies for Mild, Moderate and Severe Non-renal Lupus²

Item	Mild activity/flare BILAG C scores or single B score; SLEDAI <6	Moderate activity/flare BILAG 2 or more systems with B scores, SLEDAI 6–12	Severe activity/flare (non-renal) BILAG 1 or more A scores; SLEDAI >12
Typical manifestations attributed to lupus	Fatigue, malar rash, diffuse alopecia, mouth ulcers, arthralgia, myalgia, platelets $50\text{--}149 \times 10^9/l$	Fever, lupus-related rash up to 2/9 body surface area, cutaneous vasculitis, alopecia with scalp inflammation, arthritis, pleurisy, pericarditis, hepatitis, platelets $25\text{--}49 \times 10^9/l$	Rash involving >2/9 body surface area, myositis, severe pleurisy and/or pericarditis with effusion, ascites, enteritis, myelopathy, psychosis, acute confusion, optic neuritis, platelets $<25 \times 10^9/l$
Initial typical drugs and target doses if no contraindications	CSs ^a : topical preferred or oral prednisolone ≤ 20 mg daily for 1–2 weeks or I.m. or IA methyl-prednisolone 80–120 mg and HCQ ≤ 6.5 mg/kg/day and/or MTX 7.5–15 mg/week and/or NSAIDs (for days to few weeks only)	Prednisolone ^a ≤ 0.5 mg/day or i.v. methyl-prednisolone ≤ 250 mg $\times 1\text{--}3$ or i.m. methyl-prednisolone 80–120 mg and AZA 1.5–2.0 mg/kg/day or MTX (10–25 mg/week) or MMF (2–3 g/day) or ciclosporin ≤ 2.0 mg/kg/day and HCQ ≤ 6.5 mg/kg/day	Prednisolone ^a ≤ 0.5 mg/day and/or i.v. methyl-prednisolone 500 mg $\times 1\text{--}3$ or prednisolone $\leq 0.75\text{--}1$ mg/kg/day and AZA 2–3 mg/kg/day or MMF 2–3 g/day or CYC i.v. or ciclosporin ≤ 2.5 mg/kg/day and HCQ ≤ 6.5 mg/kg/day
Aiming for typical maintenance drugs/doses providing no contraindications	Prednisolone ^a ≤ 7.5 mg/day and HCQ 200 mg/day and/or MTX 10 mg/week Aim to reduce and stop drugs except HCQ eventually when in stable remission	Prednisolone ^a ≤ 7.5 mg/day and AZA 50–100 mg/day or MTX 10 mg/week or MMF 1 g/day or ciclosporin 50–100 mg/day and HCQ 200 mg/day; Aim to reduce and stop drugs except HCQ eventually when in stable remission	Prednisolone ^a ≤ 7.5 mg/day and MMF 1.0–1.5 g/day or AZA 50–100 mg/day or ciclosporin 50–100 mg/day and HCQ 200 mg/day; Aim to reduce and stop drugs except HCQ eventually when in stable remission

^aThe lowest effective dose of prednisolone or other CSs should be used at all times.

Appendix II: Monitoring & Patient Safety

Drug	Adverse Reactions	Monitoring	REMS
Benlysta belimumab	<ul style="list-style-type: none"> • Gastrointestinal: nausea (15%), diarrhea (12%) • Miscellaneous: infusion related reaction (17%), hypersensitivity (13%) • Pregnancy: IgG molecules cross placenta with increased amount through pregnancy (use contraception during and 4 mons. post use) 	<ul style="list-style-type: none"> • Central nervous system: worsening depression, mood changes, suicidal thought • Hypersensitivity, infusion reactions • Infections 	None needed